

A sample of X-ray emitting normal galaxies from the BMW – HRI Catalogue

Marzia Tajer^{1,2}, Ginevra Trinchieri¹, Anna Wolter¹, Sergio Campana¹, Alberto Moretti¹, Gianpiero Tagliaferri¹

¹ INAF-Osservatorio Astronomico di Brera, via Brera 28, 20121 Milano Italy

² Università di Milano - Bicocca, Dipartimento di Fisica, P.za della Scienza 3, 20126 Milano Italy

Draft: February 2, 2008

Abstract. We have obtained a sample of 143 normal galaxies with X-ray luminosity in the range $10^{38} - 10^{43}$ erg s⁻¹ from the cross-correlation of the ROSAT HRI Brera Multi-scale Wavelet (BMW – HRI) Catalogue with the Lyon-Meudon Extragalactic Database (LEDa). We find that the average X-ray properties of this sample are in good agreement with those of other samples of galaxies in the literature. We have selected a complete flux limited serendipitous sample of 32 galaxies from which we have derived the logN-logS distribution of normal galaxies in the flux range $1.1 - 110 \times 10^{-14}$ erg cm⁻² s⁻¹. The resulting distribution is consistent with the euclidean –1.5 slope. Comparisons with other samples, such as the Extended Medium Sensitivity Survey, the ROSAT All Sky Survey, the XMM - Newton/2dF survey and the *Chandra* Deep Field Survey indicate that the logN-logS distribution of normal galaxies is consistent with an euclidean slope over a flux range of about 6 decades.

Key words. X-ray – galaxies: general – X-ray surveys

1. Introduction

Detailed X-ray studies of normal galaxies have been possible only with the advent of imaging instruments aboard the *Einstein* observatory. The *Einstein* results were summarized in the catalogue and atlas published by Fabbiano et al. (1992), who constructed a large homogeneous sample of 493 galaxies, included in either A Revised Shapley-Ames Catalog of Bright Galaxies (Sandage & Tammann, 1981), or in the Second Revised Catalog of Bright Galaxies (de Vaucouleurs et al., 1976), both targets and serendipitously detected in *Einstein* fields. While representative of the galaxy population (see e.g. Shapley et al., 2001; Eskridge et al., 1995), it was not constructed to be a complete, unbiased sample, and it is likely to contain unknown selection biases.

Elliptical galaxies were found to retain large amounts ($10^8 - 10^{11}$ M_⊙) of hot gas ($T \sim 10^7$ K) whose thermal emission dominates their X-ray luminosities, while in normal spirals the integrated contribution of the evolved stellar sources, such as supernova remnants and X-ray binaries, is generally the dominant component (see Fabbiano, 1989; Fabbiano et al., 1992; Kim et al., 1992). Extended emission from a hot, gaseous component in spiral galaxies was detected only in some cases (Fabbiano & Trinchieri,

1987; Vogler & Pietsch, 1996; Trinchieri et al., 1988), or associated with starburst activity.

Subsequent observations of individual sources made by *ROSAT* and *ASCA* confirmed most of the *Einstein* results and added interesting information on the X-ray properties of normal galaxies (see among others Roberts & Warwick, 2000; Read et al., 1997; Brown & Bregman, 1998; Beuing et al., 1999) in the local universe.

With the launch of *XMM - Newton* and *Chandra*, the study of the X-ray properties of “normal” galaxies at intermediate ($z \sim 0.1$) or cosmological distances (Brandt et al., 2001; Hornschemeier et al., 2002, 2003; Georgakakis et al., 2003, 2004a,b; Norman et al., 2004) was made possible, thanks to significantly improved sensitivity, spatial and spectral resolution of the instruments. In spite of the large number of papers, however, a truly complete sample of X-ray emitting normal galaxies in the local universe with a significant number of objects has not been properly discussed in the literature so far. Georgakakis et al. (2003) give galaxy number density at $F_{0.5-2} \sim 7 \times 10^{-16}$ erg cm⁻² s⁻¹ from stacking analysis, while Georgakakis et al. (2004a) present a sample of 26 X-ray sources detected in an area of ~ 2.5 deg⁻², of which only 2 are however classified as normal galaxies. Only recently Georgakakis et al. (2004b) have presented a larger

Send offprint requests to: M. Tajer

Correspondence to: tajer@brera.mi.astro.it

sample of 11 normal galaxies detected in an area of ~ 4.5 deg $^{-2}$.

Two other samples of “normal” galaxies are available in the literature, selected in the *Chandra* Deep Fields (Hornschemeier et al., 2003; Norman et al., 2004). However their median redshifts ($z = 0.297$, Hornschemeier et al. (2003) and $z = 0$ to $z = 1.3$, Norman et al. (2004)) indicates that they should not be considered as “local”.

The large database provided by ROSAT has been exploited only marginally to derive unbiased and complete sample of galaxies. Zimmermann et al. (2001) have selected a sample of *candidate normal galaxies* from the ROSAT All Sky Survey (RASS) Bright Source Catalogue (Voges et al., 1999) above a flux limit of about 10^{-12} erg cm $^{-2}$ s $^{-1}$ (0.1 – 2.4 keV band). A few samples have been derived from the ROSAT Position Sensitive Proportional Counter (PSPC) pointed observations such as the WGA (White, Giommi, & Angelini, 1994) and the ROSPSPC (ROSAT team 2001), and a new catalogue of galaxies is in progress (G. Peres, private communication). Here we exploit the potential provided by the Brera Multi-scale Wavelet (BMW – HRI) catalogue (Panzer et al., 2003) to extract a sample of normal galaxies, as we discuss in the next sections. While the PSPC is probably more efficient at detecting faint and extended sources such as galaxies, the sharp core of the HRI point spread function allows the detection of sources in more crowded fields and to establish the extension for bright small-size sources, providing a good complement to the PSPC data.

2. The sample

In order to create a complete, serendipitous sample of galaxies with X-ray emission, we made use of X-ray data from the BMW ROSAT HRI catalogue and optical data from LEDA (Lyon-Meudon Extragalactic Database). The BMW – HRI catalogue consists of 29089 X-ray sources detected in 4303 ROSAT HRI pointed fields with exposure times longer than 100 s using a multiscale wavelet algorithm (Lazzati et al., 1999; Campana et al., 1999; Panzer et al., 2003). Sources detected with a significance $\geq 4.2 \sigma$ are contained in the catalogue, that provides name, position, count rate, flux and extension along with the relative errors. In our study we used the full catalogue, but we excluded X-ray sources in the Trapezium field, which is a rich stellar cluster in the Milky Way, where the high density of the X-ray sources would prevent proper optical identifications. The BMW – HRI catalogue can be searched via the HEASARC Browse¹ or via the Brera Observatory web site.²

¹ <http://heasarc.gsfc.nasa.gov/cgi-bin/W3Browse/w3browse.pl>

² <http://www.merate.mi.astro.it:8081/interroga/dbServer?cmd=BMW2>

Created in 1983 at Lyon Observatory, LEDA³ has been the first database of extragalactic objects and it is continuously updated. It gives a free access to the main astrophysical parameters (coordinates, morphological type, diameter and axis ratio, apparent magnitudes and colors, radial velocity, surface brightness, etc) for about 10^6 galaxies over the whole sky. The completeness in apparent B - magnitude is satisfied up to $m_B = 15.5$ (see Patirel et al., 1997).

To obtain a representative sample of galaxies we started from the BMW – HRI catalogue and included only serendipitous detections, avoiding the targets. We chose a $3'$ radius to define the typical region of the target, and selected only sources at off-axis angles $\theta > 3'$. In spite of their off-axis location, we had to exclude 12 additional sources that were targets of the observations. We then cross - correlated the positions of the X-ray sources in the BMW – HRI catalogue with those of galaxies present in the LEDA database version of 1999, with a tolerance of $20''$, which should be a reasonable guess to detect extended objects like galaxies and to avoid most of chance coincidences. This criterium is not appropriate for very extended galaxies (like e.g. M 31 or M 33), where a large number of sources is detected at distances significantly larger than our tolerance radius. Therefore we could be selecting against large size galaxies, if there is no source within a distance of $\sim 20''$ from the nucleus. However, this should not be a concern in this study, since the surface density of large galaxies is small: in particular, in the LEDA catalog, the density of galaxies with $D_{25} > 3'$ is $< 3 \times 10^{-2}$ deg $^{-2}$, which implies about < 10 in the area we surveyed.

To check the goodness of our choice, we plot in Fig. 1 the relative shifts between optical and X-ray positions. We use only point-like X-ray sources because in extended sources the association with a single optical object could be misleading (e.g. in groups and pairs). About 90% of identifications are within $13''$ (represented by the dashed circle in Fig. 1).

This is in good agreement with the HRI positional uncertainty, since the best attitude solution guarantees that on average known objects are detected within $10''$ of their catalog position, although with possible additional discrepancy mainly in declination (see the ROSAT Handbook e.g. at <http://heasarc.gsfc.nasa.gov/docs/rosat/ruh/handbook/handbook.html>). We verified that all X-ray sources with X-ray/optical off-set $> 10''$ are still within the galaxy (i.e. D_{25}). We conclude that the association of the X-ray sources with galaxies found by the cross-correlation is sound on positional grounds.

The cross-correlation yields 399 X-ray sources associated with 281 galaxies. Being constructed field-by-field, the BMW – HRI catalogue contains multiple detections of the same source. Therefore we only have 283 distinct X-

³ There exist several mirrors to access LEDA; we used the OAB one, <http://www.brera.mi.astro.it/hypercat/>




Fig. 1. Displacement between optical and X-ray positions for point-like sources in the total sample. The solid circle refers to the cross - correlation radius of 20''; the dashed one, with 13'' radius, contains about 90% of the identifications.

ray sources associated with galaxies (two distinct sources but of different extent are associated with both NGC 1399 and M 86; we list them in Table 1, but we consider only the largest one for computing fluxes and luminosities).

To check the results of the cross-correlation, we inspected HRI images for each source and obtained information from the NASA/IPAC Extragalactic Database (NED) to a) eliminate spurious coincidences [15 sources], b) eliminate AGNs [47 objects] and c) select clusters [57 objects].

- a . We eliminated a source when the obvious optical counterpart was not a LEDA galaxy, but a background or foreground object. This was checked both using NED and the X-ray/optical contour maps.
- b . We excluded objects classified in Véron-Cetty & Véron Catalogue (2001) as Seyfert 1 galaxies, QSOs, BL Lacs or AGNs. For Seyfert 2 galaxies, the X-ray emission could have a non nuclear origin, so we eliminated them when we saw from images that the X-ray emission was point-like and well centered on the nucleus, while we retained those for which we found an extended emission. We also retained starburst galaxies, LINERs and objects that we know from the literature could have a nuclear source but also non - nuclear X-ray emission (e.g. NGC 3079).
- c . Several galaxies lie inside a cluster: we inspected X-ray images and we eliminated those objects whose X-ray emission was indistinguishable from the cluster's, but we retained galaxies for which emission

clearly associated with the galaxy is detected above the cluster background (see e.g. PGC 12350 in Fig. 2). We retained galaxies in poor groups even when they are the brightest member since there is still ambiguity in the literature between emission from bright early-type galaxies and groups, which are often analyzed in the same context (see e.g. Mamon, 1992; Dell'Antonio et al., 1994; Pildis et al., 1995; Ponman et al., 1996; Mahdavi et al., 1997; Mulchaey & Zabludoff, 1998; Helsdon & Ponman, 2000; Mulchaey et al., 2003; Osmond & Ponman, 2004; Helsdon & Ponman, 2003; Jones et al., 2003).

Moreover, in 5 cases the same X-ray source was associated with two or more galaxies in a pair or in a group: since we could not discriminate on a positional basis, we chose the brightest galaxy in the pair or in the group.

These selection criteria yielded a total of 195 X-ray sources (including multiple detections) associated with 143 galaxies whose properties are listed in Table 1, available in electronic form. We report here the first page as an example. Column (1) gives the BMW – HRI name of the source, columns (2) and (3) the position of the X-ray peak, column (4) labels the source as point-like (p) or extended (e), column (5) gives the radius in which we computed count rates for extended sources (see section 3), column (6) the major semiaxis of the ellipse, for extended X-ray sources for which count rates have been computed in elliptical regions (see section 3), column (7) the minor semiaxis of the ellipse, column (8) the LEDA galaxy associated to the X-ray source, column (9) the name of the galaxy in other common catalogues (e.g. NGC), column (10) the morphological type, column (11) the distance of galaxy, column (12) the apparent B - magnitude corrected for galactic extinction, inclination and redshift effects (see Paturel et al., 1997), columns (13) and (14) the HRI count rate and error, columns (15) and (16) the X-ray flux and error (0.1 – 2 keV; count rates and fluxes have been re-computed with respect to values reported in the BMW – HRI catalogue; see section 3 for details), column (17) the logarithm of X-ray luminosity and column (18) specifies whether the galaxy is in the complete subsample (c), in cluster (Cl) or in group (Gr).

Information about magnitudes and redshifts are from LEDA and NED except for 3 objects that we observed ourselves (see section 5). We calculated distances from redshifts, assuming $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$; when the heliocentric radial velocity of galaxy was less than 3000 km s^{-1} , however, we used distances from Nearby Galaxy Catalogue (Tully, 1988) corrected for $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

One hundred and sixteen of the 143 galaxies have redshift and magnitude (including our observations). Nineteen galaxies have m_B and no z , 3 have z and no m_B and 5 have neither z nor m_B . The redshift distribution ranges from $z = 0$ from $z \sim 0.15$, but $\sim 90\%$ of galaxies have $z < 0.07$.

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
BMW000523.9+161307	00 05 23.83	+16 13 11.3	e	52				PGC 0000372		698		15.74	2.49	7.40	1.17	43.59	Cl
BMW002055.1+215208	00 20 55.43	+21 51 52.2	p					PGC 0001333 IC 1543	Sbc	112	14.54	7.63	1.11	3.58	0.52	41.71	c
BMW002055.2+215208																	
BMW002549.3-453227	00 25 49.26	-45 32 26.5	p					PGC 0143535	Sab		16.57	18.64	3.16	7.64	1.30		
BMW002950.2-405630	00 29 50.99	-40 56 37.9	p					PGC 0130966 DUKST 294-9	Sc	241	15.35	19.06	2.31	7.81	0.95	42.72	c
BMW003652.3-333310	00 36 52.82	-33 33 14.7	p					PGC 0002204 ESO 350-IG38	S?	123	14.96	2.74	0.43	1.12	0.14	41.27	c
BMW003918.5+030220	00 39 18.47	+03 02 14.8	p					PGC 0002362 NGC 194	E	103	13.09	14.75	1.94	6.49	0.85	41.89	c; Gr
BMW003948.3+032219	00 39 48.77	+03 22 21.0	p					PGC 0002401 UM 57				6.15	0.49	2.70	0.22		
BMW004242.0+405154	00 42 42.48	+40 51 52.6	e	60				PGC 0002555 NGC 221	E	1	8.18	52.44	1.06	26.75	0.54	38.52	Gr
BMW010716.2+323117	01 07 16.20	+32 31 16.6	p					PGC 0003966 NGC 379	S0	110	13.48	1.37	0.35	0.67	0.17	40.98	Gr
BMW010717.4+322857	01 07 17.76	+32 28 57.0	p					PGC 0003969 NGC 380	E	88	13.24	4.19	0.50	2.06	0.24	41.26	Gr
BMW010717.8+322857																	

- (1) BMW – HRI source name
(2) X-ray coordinates: RA (J2000)
(3) X-ray coordinates: Dec (J2000)
(4) X-ray source point-like (p) or extended (e)
(5) Radius of circle in which count rates were computed, for extended X-ray sources
(6) Major semiaxis of ellipse in which count rates were computed, for some peculiar extended X-ray sources
(7) Minor semiaxis of ellipse in which count rates were computed, for some peculiar extended X-ray sources
(8) LEDA galaxy associated
(9) Other galaxy name
(10) Morphological galaxy type
(11) Galaxy distance (Mpc)
(12) Apparent B - magnitude
(13) X-ray count rate (10^{-3} count s^{-1})
(14) X-ray count rate error (10^{-3} count s^{-1})
(15) X-ray flux (0.1 – 2 keV band; 10^{-13} erg cm^{-2} s^{-1})
(16) X-ray flux error (0.1 – 2 keV band; 10^{-13} erg cm^{-2} s^{-1})
(17) Logarithm of X-ray luminosity (0.1 – 2 keV band; erg s^{-1})
(18) Notes: c if the galaxy is in the complete subsample, Cl if in cluster, Gr if in group

Table 1. First page of Table of the total sample, available in electronic form.

2.1. The complete serendipitous subsample

To study the general properties of the sample we need to derive a subsample with well known completeness criteria and limits. To this end we constructed a complete sample with both X-ray and optical flux limits. We must consider both the X-ray and the optical completeness criteria. The X-ray completeness is related to the BMW – HRI catalogue, that includes all sources with a significance $\geq 4.2 \sigma$. To take into account the optical limits, we excluded galaxies fainter than $m_B = 15.5$ (the completeness limit assumed for the LEDA Catalogue, see Paturel et al., 1997).

We have also excluded objects at low galactic latitude ($|b| \leq 10^\circ$) to avoid source confusion in the galactic plane and X-ray sources with off-axis angle $\theta \geq 18'$ to match the circular HRI field of view used in the sky coverage computation (see section 7).

The resulting sample of 96 objects is complete both in X-ray and optical, at the given limits. However, since galaxies are often in agglomerates, some of them are expected to be related to the targets and therefore not truly serendipitous. We therefore excluded all sources known to be associated with the target (e.g. galaxies in pairs, groups or clusters; 52 objects). When an association was not documented (e.g. from NED, LEDA) we conservatively excluded galaxies at the same redshift of the target (12 objects).

The complete, serendipitous sample of 32 galaxies thus obtained is given in Table 3 and will be used to calculate the logN-logS distribution in the local universe ($z < 0.07$), as will be described in detail in section 7.

3. X-ray characterization

The BMW – HRI catalogue provides count rates derived using the wavelet algorithm, in an automated way and under particular assumptions (for details, see Lazzati et al., 1999; Campana et al., 1999). We verified that, while count rates are correctly computed for point-like sources, for very extended sources such as NGC 1399, the extension and count rates given by the algorithm are underestimated. Moreover, when multiple observations are available we can improve the statistics by considering the full set of data. We have therefore recalculated all count rates, using the BMW – HRI positions and original HRI data retrieved from the public archives⁴, in pulse height analyzer (PHA) channels 1 – 10 to increase the signal-to-noise ratio (for a justification of this choice of PHA, see Trinchieri et al., 1997). When multiple observations of the same field are available, we summed the data if they have the same pointing coordinates and comparable exposure times. Otherwise, we typically used the longer exposures or those where the source is closer to the field’s center.

⁴ <http://wave.xray.mpe.mpg.de/>
<http://heasarc.gsfc.nasa.gov/>

and

3.1. Count rates

We classified sources as point-like or extended, based on the radial distribution of the emission relative to the shape of the HRI point spread function (PSF; for a description of the *ROSAT* PSF see Boese, 2000) at the corresponding off-axis angle. For the “extended sources” the counts are taken from the largest region that contains source counts, generally a circle of radius reported in Table 1, evaluated from the radial profile. In few cases, given the particular shape of the source, we used ellipses to evaluate the counts; we give in Table 1 minor and major semiaxis. There are also a few extended sources for which ad hoc regions (neither circles nor ellipses) have been used to estimate counts: these are PGC 5323, PGC 5324, PGC 13418, PGC 13433 and PGC 56962.

For point-like sources the counts are obtained in a circular region centered at the peak of X-ray emission, with radius that includes about 90% of source counts according to the PSF. The PSF degrades as the angular distance from the center of field increases, so we chose a radius of $r = 18''$ for $3' < \theta \leq 10'$, $r = 25''$ for $10' < \theta \leq 15'$ and $r = 40''$ for $\theta > 15'$, following Boese (2000). We evaluated the background in an annulus concentric to the source radius, with radii depending on the off-axis. When the source was particularly faint, we calculated count rates in a circle of radius corresponding to a smaller fraction of the PSF, to increase the signal-to-noise ratio. We then corrected the count rate accordingly, following Boese (2000).

We compared count rates obtained in this way with those reported in the BMW – HRI catalogue and we found a general agreement, with the exception of some sources whose extension had been largely underestimated by the wavelet algorithm, as stated above. Indeed, our count rates would be the equivalent to the “counted count rates” reported in the catalogue, rather than those computed with the wavelet algorithm. In the comparison with this quantity we found a systematically higher count rate, consistent with the larger PHA interval used (1 – 10 in our analysis and 2 – 9 in the BMW – HRI catalogue).

The resulting net count rates are given in Table 1, corrected for vignetting and lost counts due to the PSF (for point sources only).

3.2. Fluxes and luminosities

The count rates were converted into 0.1 – 2 keV fluxes using a conversion factor corresponding to a bremsstrahlung spectrum with $kT = 5$ keV plus the line of sight absorption appropriate for each source from Dickey & Lockman (1990) (reported in Table 2). Although this spectrum might not be suitable for all kinds of sources, the flux in the *ROSAT* energy window depends only weakly on the spectral model assumed, while it is more dependent on low energy absorption. The resulting fluxes are in the range $10^{-14} - 10^{-11}$ erg cm⁻² s⁻¹.

N_{H} (cm^{-2})	CF ($\text{erg cm}^{-2} \text{count}^{-1}$)
1×10^{20}	3.7×10^{-11}
2×10^{20}	4.1×10^{-11}
5×10^{20}	4.9×10^{-11}
8×10^{20}	5.5×10^{-11}
3×10^{21}	9.3×10^{-11}

Table 2. Galactic N_{H} and corresponding unabsorbed flux in the band 0.1 - 2 keV for 1 count/s, assuming a thermal bremsstrahlung spectral model with $kT = 5$ keV.

To calculate luminosities, we used distances listed in Table 1. The corresponding range in L_X is $10^{38} - 10^{43} \text{ erg s}^{-1}$.

4. The atlas

We provide in Fig. 2 an overlay of X-ray contours of the detected galaxies onto optical images from the Digital Sky Survey II (DSS II) available from the ESO archive⁵. The Figure is available in electronic edition. We report here the first page as an example. When available, we used optical images obtained with the blue filter, otherwise we used those obtained in the red filter. For PGC 209730 only the DSS I plate is available.

Galaxies, ordered in RA and generally at the center of the field, are identified by their PGC name.

X-ray contours are produced from images in the PHA range 1 – 10, smoothed with a Gaussian function with $\sigma = 5''$ for point-like sources and with $\sigma = 10''$ for extended sources.

5. Optical observations

In order to measure redshifts and magnitudes for some of the galaxies of our sample, we made spectroscopical and photometrical observations at the 1.52 meter telescope of the Osservatorio Astronomico di Bologna, at Loiano (Italy) on the nights of the 16th and 17th October 2001. Because of bad atmospheric conditions, we were able to observe only 3 galaxies. A spectrophotometric calibration star was also observed. We present the results obtained in Appendix A.

6. Comparison with literature results

To verify whether the total sample of 143 objects is representative of the X-ray properties of normal galaxies, we calculated X-ray luminosities where possible (Table 1) and we plotted the distributions of L_X , L_B , the ratio L_X/L_B and the L_X - L_B relationship for spiral and early-type galaxies. For the 19 galaxies for which the redshift is not known but we have m_B , the ratio is calculated from




Fig. 3. X-ray - to - optical ratio distribution for the total sample. Both fluxes are in units of $\text{erg cm}^{-2} \text{s}^{-1}$.

fluxes. We compared our results with those in the literature and found a good general agreement.

In particular:

1. the bulk of the galaxies in the sample has an X-ray luminosity between $10^{38} \text{ erg s}^{-1}$ and few $10^{42} \text{ erg s}^{-1}$, in accordance with, e.g., Fabbiano (1989); 7 objects have $L_X \gtrsim 10^{43} \text{ erg s}^{-1}$, but most of them lie in a group, so that the intergalactic medium could contribute to their luminosity; we cannot exclude the presence of an unidentified AGN for some of these objects.
2. For spiral galaxies we found a linear relationship between X-ray and optical luminosities ($L_X \propto L_B^{1.0 \pm 0.2}$), in agreement with Fabbiano et al. (1992). In a subsequent, more complete statistical analysis of the 234 “normal” spiral and irregular galaxies reported by Fabbiano et al. (1992), Shapley et al. (2001) and Fabbiano & Shapley (2002) find however that the $L_X - L_B$ relationship is significantly steeper than linear, with a slope of about 1.5. Our study is based on a much smaller sample (32 spiral and irregular galaxies), and small statistics could account for the discrepancy, as pointed out by Fabbiano & Shapley (2002) in comparing their findings with previous works.
3. For early-type galaxies we found a steeper relationship, $L_X \propto L_B^{1.6 \pm 0.2}$, consistent with those obtained by Fabbiano et al. (1992) and Eskridge et al. (1995).
4. Values of the X-ray - to optical ratio in our sample cover roughly the same range ($-4 < \log(L_X/L_B) < 0$ with L_X and L_B in erg s^{-1}) as the spiral galaxies in Shapley et al. (2001) and the early-type galaxies in Eskridge et al. (1995), but their distributions are different. We plot in Fig. 3 the histogram of the X-ray - to

⁵ <http://archive.eso.org/dss/dss>




Fig. 4. Sky coverage for the fields at galactic latitude $|b| \geq 10^\circ$ and off-axis angle $3' \leq \theta \leq 18'$, computed by assuming a thermal bremsstrahlung spectrum with $kT = 5$ keV and galactic line of sight absorption.

- optical ratio of the total sample of 143 galaxies. We will analyze this subject in greater detail in subsection 7.1.

7. The logN-logS distribution

The complete serendipitous sample of galaxies derived in section 2.1 (listed in Table 3) was used to calculate the integral flux distribution (logN-logS) of normal galaxies with X-ray emission above the X-ray flux limit of the BMW – HRI Catalogue and B - magnitude ≤ 15.5 .

The sensitivity of the HRI instrument is not uniform over the entire field of view. Moreover the observing time is different for different fields so we need to calculate the area surveyed at any given flux (sky coverage).

In the BMW – HRI catalogue the published sky coverage was calculated by means of simulations (see Panzera et al., 2003). In this work we used a sky coverage calculated with the same procedure, but with parameters that reflect our source selection criteria. Therefore we included only fields with galactic latitude $|b| \geq 10^\circ$, we considered only an annular region with $3' \leq \theta \leq 18'$ (the lower limit is to account for the target region; the upper limit is the largest radius within the field of view of the detector in the assumption of circular symmetry) and we assumed a bremsstrahlung spectrum with $kT = 5$ keV plus the line of sight absorption.

The resulting sky coverage is plotted in Fig. 4. The maximum area is $\sim 314 \text{ deg}^2$ and corresponds to fluxes above $\sim 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$. The surveyed area is $\sim 196 \text{ deg}^2$ at $10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ and $\sim 3 \text{ deg}^2$ at $\sim 1.1 \times$




Fig. 5. The integral logN-logS distribution for the complete serendipitous subsample (asterisks); the solid line represents the -1.1 slope and the dashed line the euclidean -1.5 slope.

$10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ (the lowest flux for the galaxies in our sample).

For consistency with the sky coverage calculation, the fluxes given in Table 3 are derived from the original count rates estimated in the BMW – HRI catalogue using the wavelet algorithm and PHA channels of HRI from 2 to 9, which are corrected for vignetting and PSF according to Campana et al. (1999) and computed for the energy band $0.5 - 2 \text{ keV}$, the standard of BMW – HRI Catalogue.

The integral logN-logS distribution of the sample is shown in Fig. 5 and covers two decades in flux, from ~ 1.1 to $\sim 110 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$. The overall distribution could be approximated with a slope of ~ -1.1 (solid line in Fig. 5). However the euclidean slope of -1.5 (dashed line) is also consistent with the data: the excess of galaxies at the highest fluxes is small and consistent within the limited statistics. Moreover, at the lower fluxes, we could have some problems with incompleteness, as we discuss below.

7.1. Comparison with the literature: X-ray - to - optical ratio distribution

Before comparing our logN-logS distribution with those of other samples in the literature, we need to better investigate the X-ray - to - optical ratio distribution of our complete sample. We will consider samples derived from *ROSAT* and *Einstein* observations that cover a flux range similar to ours. We will include samples derived from *XMM - Newton* and *Chandra* surveys, that cover a flux

LEDA name	N_H (10^{20} cm^{-2})	Count rate ($10^{-3} \text{ count s}^{-1}$)			$F_{0.5-2}$ ($10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$)		
0157406	4.9	0.37	±	0.07	0.12	±	0.02
0068536	2.5	1.11	±	0.34	0.32	±	0.09
0025825	4.2	1.12	±	0.22	0.35	±	0.06
0042833	2.0	1.31	±	0.23	0.35	±	0.05
0028990	0.8	1.50	±	0.30	0.36	±	0.07
0005323	3.8	1.33	±	0.28	0.41	±	0.09
0010446	3.1	1.48	±	0.24	0.43	±	0.06
0057078	2.3	1.63	±	0.28	0.44	±	0.08
0069338	8.8	1.24	±	0.20	0.46	±	0.07
0038773	2.0	1.70	±	0.11	0.46	±	0.03
0001333	4.2	1.55	±	0.27	0.48	±	0.09
0007289	5.5	1.71	±	0.26	0.56	±	0.10
0070861	2.6	2.11	±	0.38	0.61	±	0.12
0018991	5.4	2.18	±	0.32	0.70	±	0.10
0004117	4.0	2.36	±	0.23	0.73	±	0.06
0043675	3.9	2.49	±	0.37	0.77	±	0.12
0002204	1.9	3.23	±	0.23	0.87	±	0.05
0045318	1.0	3.78	±	0.60	0.91	±	0.14
0046432	3.2	3.13	±	0.66	0.91	±	0.20
0047432	2.3	3.61	±	0.64	0.97	±	0.16
0130966	2.4	4.41	±	0.81	1.19	±	0.22
0016574	5.9	3.73	±	0.51	1.23	±	0.17
0005324	3.8	4.34	±	0.51	1.35	±	0.16
0057728	1.5	5.11	±	0.39	1.38	±	0.11
0013368	1.4	7.22	±	1.41	1.73	±	0.24
0029050	0.8	10.20	±	0.70	2.45	±	0.17
0002362	2.7	9.06	±	0.99	2.63	±	0.29
0004848	6.1	9.26	±	2.43	3.06	±	0.66
0063122	7.9	11.02	±	1.42	3.97	±	0.36
0006367	3.0	14.80	±	0.60	4.29	±	0.17
0017451	7.5	15.66	±	1.72	5.64	±	0.72
0028995	4.8	34.20	±	2.00	10.94	±	0.64

Table 3. Galactic column density (from Dickey & Lockman 1990), BMW count rates and fluxes for galaxies in the complete serendipitous sample, ordered by increasing flux.

range significantly fainter than ours, in the discussion of the logN-logS (subsection 7.2).

The best available comparison could be with the sample of *candidate normal galaxies* found by Zimmermann et al. (2001) in the *ROSAT* All Sky Survey (RASS) and with the normal galaxies found in the Einstein Extended Medium Sensitivity Survey (EMSS; Gioia et al., 1990), both X-ray selected. An effectively optically selected sample for comparison is the *Einstein* galaxy sample (Fabbiano et al., 1992; Shapley et al., 2001; Eskridge et al., 1995).

Zimmermann et al. (2001) made a correlation study of the RASS Bright Source Catalogue (Voges et al., 1999) with the Catalogue of Principal Galaxies (Paturel et al., 1989), from which they selected a sample of 198 *candidate galaxies*, i.e. X-ray sources whose optical counterpart was not designated as AGN in the literature. These selection criteria are similar to ours, and the Catalogue of Principal Galaxies is a preliminary version of the current LEDA database, so the two samples can be easily compared; how-

ever most of the Zimmermann et al. (2001) sources have fluxes above $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ (computed in the 0.1–2.4 keV band, assuming a power law spectrum with photon index $\Gamma = 2.3$).

The EMSS was obtained from the analysis of 1453 images of the imaging proportional counter (IPC) on board the *Einstein* Observatory. The survey covers an area of 778 deg^2 at $|b| > 20^\circ$ with limiting sensitivity ranging from $\sim 5 \times 10^{-14}$ to $\sim 3 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.3–3.5 keV band). 835 serendipitous sources were detected at or above the 4σ level (see Gioia et al., 1990; Stocke et al., 1991). Among these, 17 were identified as normal galaxies.

The *Einstein* sample is the catalogue of normal galaxies observed by the *Einstein* satellite, compiled by Fabbiano et al. (1992) and reanalyzed by Eskridge et al. (1995, for early-type galaxies) and by Shapley et al. (2001, for spiral galaxies); we no longer distinguish here the two samples since early and late type galaxies are mixed in the BMW – HRI sample.




Fig. 6. B flux versus X-ray flux for our total sample (solid triangles), for the *Einstein* sample (empty circles), for the Zimmermann et al. 2001 sample galaxies (asterisks), and for the EMSS galaxies (empty pentagons). Solid lines correspond to $\log(F_X/F_B) = 0, -2, -4$, as indicated; the horizontal dashed line is the $m_B = 15.5$ limit. We have considered only detections in the *Einstein* sample: the exclusion of upper limits should not affect the results, since their distribution is consistent with that of the detections (see Fig. 5 in Shapley et al. 2001). We do not plot objects at $F_{0.1-2} < 2 \times 10^{-12}$ erg cm $^{-2}$ s $^{-1}$ in the Zimmermann et al. (2001) sample because below this flux their sample is not complete (see text).

In Fig. 6 we plot the distribution of F_X and F_B values from all samples considered. For consistency with the values in Table 1, all fluxes are converted to the 0.1 – 2 keV energy band, using a thermal bremsstrahlung spectrum with temperature $kT = 5$ keV and $N_H = 3 \times 10^{20}$ cm $^{-2}$.

Fig. 6 clearly indicates that galaxies belonging to different samples populate different regions in the plot. Our sample (solid triangles) has an X-ray flux range between 10^{-14} and 10^{-11} erg cm $^{-2}$ s $^{-1}$ and $\log(F_X/F_B)$ between -4 and 0 (see also Fig. 3). The bulk of *Einstein* galaxies (empty circles) is typically at higher average fluxes and at lower values of $\log(F_X/F_B)$ (between -4 and -2) compared to our distribution and the X-ray selected samples in general. This sample is the largest, it is effectively optically selected and reasonably clean of the contamination from AGN (Shapley et al., 2001). However, since it is not complete, it might not provide the true distribution of the X-ray - to - optical ratios.

The second largest sample is derived from Zimmermann et al. (2001), with an additional flux limit $F_X > 2 \times 10^{-12}$ erg cm $^{-2}$ s $^{-1}$ (for completeness,

see discussion below) and an upper limit in luminosity at $\log L_X = 42.7$ erg s $^{-1}$ (the highest luminosity in our complete sample), comparable to their limit to exclude potential AGNs from the sample. The distribution of *candidate galaxies* of Zimmermann et al. (2001) (asterisks) is significantly different from that of the *Einstein* sample and extends at $\log(F_X/F_B) > 0$.

The F_X/F_B distribution for EMSS galaxies (empty pentagons) is at intermediate values and more consistent with that of our sample.

Also plotted in Fig. 6 is the optical flux limit applied to our sample, $m_B = 15.5$. It is evident that the exclusion of galaxies fainter than this for the logN-logS calculation has an effect that increases as flux decreases. We have attempted to quantify it in order to correct the curve for lost objects. Unfortunately we could not properly estimate the correction because none of the galaxy samples available allows us to derive the true distribution of X-ray - to - optical ratios of normal galaxies. The EMSS should represent the true F_X/F_B distribution, but, given the small number of galaxies (17, of which only 15 with B-magnitude), statistical errors are large. The difference in the distribution of ratios between the two larger samples in Fig. 6 suggests that they might be affected by opposite biases: the Zimmermann et al. (2001) sample is likely to contain unclassified AGNs, while the *Einstein* sample could lose objects at the highest X-ray - to - optical ratios. If we use the three samples to estimate how many galaxies are lost as a function of X-ray flux, we find that the corrections to the logN-logS are small and the recomputed curve is consistent with the euclidean slope in the observed flux range.

We also tried to estimate the correction by considering optically fainter galaxies. At $m_B = 16$, LEDA is about 90% complete (see Fig. 7 in Paturel et al., 1997): if we include galaxies down to this flux limit, we only add 4 objects to our serendipitous sample, distributed over the whole range of X-ray fluxes. Their inclusion only has a marginal influence on the normalization, and none on the slope of the distribution.

We conclude that, since the effects introduced by the optical limit are small, the logN-logS we derive is consistent with the euclidean slope.

7.2. Comparison with the literature: logN-logS

Figure 7 shows the comparison between the logN-logS derived above with several available from the literature. All fluxes are recomputed in the 0.5 – 2 keV range that we use.

We find an excellent agreement with other samples that cover similar or brighter flux ranges than the present sample.

The *candidate galaxies* (Zimmermann et al., 2001) appear to connect smoothly with the euclidean extrapolation of the BMW – HRI logN-logS above 10^{-12} erg cm $^{-2}$ s $^{-1}$. We interpret the flattening observed in the




Fig. 7. Comparison of the logN-logS distribution obtained from the present sample (asterisks) with several other relations for normal galaxies from the literature. The dashed line represents the euclidean relationship of Figure 5 extrapolated to the full flux range considered. Symbols refer to: EMSS galaxies (empty circles); *candidate galaxies* in Zimmermann et al. 2001 (solid squares); normal galaxies (crosses), including LLAGN (solid circles) (Georgakakis et al. 2004a), and stacking analysis (solid triangles, Georgakakis et al. 2003), found in the *XMM-Newton*/2dF survey; *XMM-Newton*/Sloan DSS normal galaxies from Georgakakis et al. (2004b) (empty squares); *Chandra* Deep Fields results from Hornschemeier et al. (2003) (empty triangles); optimistic or pessimistic curves given by Bauer et al. (2004) (upper/lower dotted line); Bayesian sample of Norman et al. (2004) reported by Ranalli et al. (2004) (dot – dashed line). Open stars and horned-shaped box indicate the detections and fluctuation analysis predictions of Miyaji & Griffiths (2002).

Zimmermann et al. (2001) data at lower fluxes as a result of their selection criteria. In fact, they indicate a $\geq 90\%$ completeness for count rates ≥ 0.1 count s^{-1} , which converts to a flux of $F_{0.5-2} \sim 7 \times 10^{-13}$ erg cm^{-2} s^{-1} .

Although small, the EMSS is a truly complete sample, since it is serendipitously X-ray selected and virtually completely identified (Gioia et al., 1990; Stocke et al., 1991; Maccacaro et al., 1994). The EMSS appears to be euclidean and almost coincident with our curve for fluxes above 10^{-13} erg cm^{-2} s^{-1} .

We have also extended the comparison to include samples at fainter fluxes. Georgakakis et al. (2004a) have computed the logN-logS of sources in the *XMM-Newton*/2dF survey, obtained with the EPIC instrument on board the *XMM-Newton* satellite. This survey covers an area of about 2.5 deg 2 to the flux limit $\sim 10^{-14}$ erg

cm^{-2} s^{-1} in the $0.5 - 8$ keV band (or $F_{0.5-2} \sim 5 \times 10^{-15}$ erg cm^{-2} s^{-1}). They find two “normal” galaxies in their sample, that imply a density of < 1 source at $F_X \sim 10^{-14}$ erg cm^{-2} s^{-1} , lower than what we find.

However, the sample of Georgakakis et al. (2004a) contains three additional galaxies (at $z \leq 0.1$ with $L_{0.5-8} \sim 10^{42}$ erg s^{-1}) that the authors do not consider because they might contain Low Luminosity AGNs (LLAGNs, see their Fig. 2). Since we cannot exclude that our sample also contains a few LLAGNs (see above and section 6), we should consider these objects for a better comparison with our sample. With the inclusion of these objects the logN-logS better matches ours.

Also shown in Fig. 7 are the constraints from the stacking analysis results of Georgakakis et al. (2003) at fainter fluxes, computed from optically selected galaxies at a mean redshift of $z \sim 0.1$. The point derived from the total sample considered is in excellent agreement with our logN-logS.

Recently Georgakakis et al. (2004b) have presented a pilot sample of normal galaxies serendipitously detected in *XMM-Newton* public observations. They found 11 “normal” galaxy candidates with luminosities below 10^{42} erg s^{-1} over an area of ~ 4.5 deg 2 . They find that the LogN-LogS derived from this sample (plotted in Fig. 7) is again almost euclidean in slope, although at a smaller normalization than ours.

We also plot the results from deeper surveys, using data from Hornschemeier et al. (2003), Norman et al. (2004) as reported by Ranalli et al. (2004), Bauer et al. (2004), derived from the *Chandra* Deep Fields. All these relations fall close to, although in general below, the extrapolation from our sample. Different authors derive different slopes for their samples, but they are all consistent with the euclidean one.

The lower normalizations found in these latter samples could be explained in part by the combined effects of more stringent criteria to minimize contamination from the AGNs, even though of low luminosity, and of different relative occurrence of the galaxy types (spiral/starburst vs early types).

As already discussed, some residual contamination from low luminosity AGNs could be present in the sample we have considered, since we have little information on the optical spectra, and could only reject known AGNs. We note however that Zimmermann et al. (2001) have applied the same criterium adopted by Georgakakis et al. (2004a), namely an X-ray to optical luminosity ratio smaller than 10^{-2} , and that the EMSS sample, which is well studied optically, should not be contaminated by AGNs. We have nevertheless considered discarding sources in our sample that have a $\log(F_{0.5-2}/F_B) < -2$. We have 7 objects that violate this limit, mostly at the high flux end. The resulting logN-logS relation is slightly steeper, but consistent with that presented in Fig. 5, and would not significantly lower the normalization of the “euclidean” curve plotted. However the location of the different points from Georgakakis et al. (2004a) that consider/discard possible

contamination from LLAGNs give an idea of the possible uncertainties involved.

The effect of different relative contributions from the early/late types is more complicated to assess. The lowest flux points (Hornschemeier et al., 2003) are derived from late type galaxies, so they could underrepresent the total population. However, Bauer et al. (2004) suggest that the early type galaxies follow a flatter distribution, and in any case their LogN-logS is always below those from starburst/quiescent galaxies (see their Fig. 9), so their contribution could in fact be negligible at the fainter fluxes. In the range covered by Georgakakis et al. (2004b), there is only one early type galaxy (but the sample is very small), while in our sample we have a sizeable fraction of early types (1/3 among the objects with a morphological classification), and the percentage increases to $\sim 50\%$ in the “candidate normal galaxies” of Zimmermann et al. (2001), for which moreover there is no apparent significant difference in the two slopes either. The stacking analysis results from Georgakakis et al. (2003) derived separately for E/S0 and Sa-Scd bracket the extrapolation of the euclidean logN-logS obtained from brighter samples, with Sa-Scd in better agreement with the samples at low fluxes. Since the emission from early and late type galaxies is due to significantly different processes (Fabbiano, 1989; Fabbiano et al., 1992; Kim et al., 1992; Eskridge et al., 1995; Shapley et al., 2001), a different evolution is not out of the question. An assessment of the local LogN-logS separately for the different morphological types would be a step forward to a better understanding of the properties of galaxies as a class, and would provide stronger constraints for the investigation of normal galaxies at higher redshifts. Investigating this aspect is beyond the scope and the potential of this work. For the present time, we simply notice how remarkable it is that, in spite of the different selection criteria and instruments used to define all the samples considered, the surface densities of normal galaxies is consistent with a single euclidean distribution for about 6 decades in flux (from $\sim 10^{-11}$ to $\sim 3 \times 10^{-17}$ erg cm $^{-2}$ s $^{-1}$).

8. Conclusions

We present the results for an “almost serendipitous” sample of 143 X-ray emitting normal galaxies selected from the cross-correlation of the BMW – HRI Catalogue and the LEDA database. Isointensity X-ray contours are overlaid onto the optical images for all galaxies and are presented in an atlas in Fig. 2. The X-ray characteristics of the sample, listed in Table 1, are derived in a uniform way and are used in comparison with other samples in the literature. We find that the general properties of the total sample are in good agreement with those already known for normal galaxies.

We have also selected a complete subsample of 32 truly serendipitous sources in the local universe ($z < 0.07$), for which we derived the logN-logS distribution in the flux range between ~ 1.1 and 110×10^{-14} erg cm $^{-2}$ s $^{-1}$, in

the 0.5 – 2 keV energy band. We find that the relation is consistent with the euclidean distribution.

Moreover, we find a good agreement between our logN-logS and those derived from *ROSAT* PSPC and *Einstein* data at similar or brighter fluxes and from *XMM - Newton* and *Chandra* at fainter fluxes: the overall distribution appears to be consistent with an euclidean slope for about 6 decades in flux, from $\sim 3 \times 10^{-17}$ to 10^{-11} erg cm $^{-2}$ s $^{-1}$. The normalizations of different samples are consistent within a factor of ~ 2 .

Although with limited statistics, this work provides a first estimate of the number density of sources identified with normal galaxies in the nearby universe, in a flux range (both optical and X-ray) easily accessible for follow up detailed observations. This will allow us to provide a solid basis that will help in studying and classifying objects found in deeper surveys.

While current efforts are mainly focused on probing the distant universe to determine the relevance of normal galaxies as a class at very faint fluxes, the success of these studies depends also on the constraints given by the bright flux end of the number counts, which is still poorly studied. This is particularly relevant at the lowest fluxes where the number counts approach the total source density (e.g. the detections and fluctuation analysis results of Miyaji & Griffiths (2002), see Hornschemeier et al. (2002) and Fig. 7). The sample here derived will be therefore instrumental to studies of both the galaxy cosmological evolution and the contribution of this class of sources to the X-ray background.

Acknowledgements. We thank Dr. R. Panzera for her invaluable help, in particular in the extraction of our sample from the BMW – HRI catalogue.

We thank Dr. U. Zimmermann for useful discussion on their work and for providing us with the original data for their logN-logS that we use in Fig. 7.

We acknowledge partial financial support from the Italian Space Agency (ASI).

This research has made use of the Lyon-Meudon Extragalactic Database (LED A), <http://www-obs.univ-lyon1.fr/hypercat/> or <http://www.brera.mi.astro.it/hypercat/>.

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

This research has made use of the SAOimage ds9 developed by the Smithsonian Astrophysical Observatory.

The compressed files of the “Palomar Observatory - Space Telescope Science Institute Digital Sky Survey” of the northern sky, based on scans of the Second Palomar Sky Survey are copyright (c) 1993-2000 by the California Institute of Technology and are distributed by agreement. All Rights Reserved.

References

Bauer, F. E., Alexander, D. M., Brandt, W. N. et al., 2004, accepted for publication in *AJ*, astro-ph/0408001

- Beuing, J., Döbereiner, S., Böhringer, H. & Bender, R. 1999, MNRAS 302, 209
- Boese, F. G. 2000, A&AS 141, 507
- Brandt, W. N., Hornschemeier, A. E., Alexander, D. M., et al. 2001, AJ 122, 1
- Brown, B. A. & Bregman, J. N. 1998, ApJ 495L, 75
- Campana, S., Lazzati, D., Panzera, M. R. & Tagliaferri, G. 1999, ApJ 524, 423
- David, L. P., Jones, C. & Forman, W., 1992, ApJ 388, 82
- Dell’Antonio, I. P., Geller, M. J. & Fabricant, D. G. 1994, AJ 107 427
- Dickey, J. M. & Lockman, F. J. 1990, ARA&A 28, 215
- Eskridge, P. B., Fabbiano, G. & Kim, D.-W. 1995, ApJS 97, 141
- Fabbiano, G. 1989, ARA&A 27, 87
- Fabbiano, G. & Shapley, A., 2002, ApJ 565, 908
- Fabbiano, G. & Trinchieri, G. 1987, ApJ 315, 46
- Fabbiano, G., Kim, D.-W. & Trinchieri, G. 1992, ApJS 80, 531
- Georgakakis, A., Georgantopoulos, I., Stewart, G. C., Shanks, T., & Boyle, B. J. 2003, MNRAS 344, 161
- Georgakakis, A., Georgantopoulos, I., Vallbe, M., et al. 2004a, MNRAS 349, 135
- Georgakakis, A., Georgantopoulos, I., Basilakos, S., Plionis, M. & Kolokotronis, V., 2004b, MNRAS in press, astro-ph/0407387
- Gioia, I. M., Maccaro, T., Schild, R. E., et al. 1990, ApJS 72, 567
- Helsdon, S. F. & Ponman, T. J. 2000, MNRAS 319, 933
- Helsdon, S. F. & Ponman, T. J. 2003, MNRAS 340, 485
- Hornschemeier, A. E., Bauer, F. E., Alexander, D. M., et al. 2003, AJ 126, 575
- Hornschemeier, A. E., Brandt, W. N., Alexander, D. M., et al. 2002, ApJ, 568, 82
- Jones, L. R., Ponman, T. J., Horton, A., et al. 2003, MNRAS 343, 627
- Kim, D.-W., Fabbiano, G., & Trinchieri, G. 1992, ApJS, 80, 645
- Lazzati, D., Campana, S., Rosati, P., Panzera, M. R. & Tagliaferri, G. 1999, ApJ 524, 414
- Maccaro, T., Gioia, I. M., Wolter, A., Zamorani, G. & Stocke, J. T. 1988, ApJ 326, 68
- Maccaro, T., Wolter, A., McLean, B., et al. 1994, ApL&C 29, 267
- Mahdavi, A., Böhringer, H., Geller, M. J. & Ramella, M. 1997, ApJ 483, 68
- Mamon, G. A. 1992, in *Physics of Nearby Galaxies: Nature or Nurture?*, ed T. X. Thuan, C. Balkowski & J. Tran Thanh Van (Gif-sur-Yvette:Editions Frontières), 367
- Miyaji, T. & Griffiths, R. E., 2002 ApJ 564L, 5
- Mulchaey, J. S. & Zabludoff, A. I. 1998, ApJ 496, 73
- Mulchaey, J. S., Davis, D. S., Mushotzky, R. F. & Burstein, D., 2003, ApJS 145, 39
- Norman, C., Ptak, A., Hornschemeier, A., et al. 2004, ApJ 607, 721
- Osmond, J. P. F. & Ponman, T. J., 2004, MNRAS 350, 1511
- Panzera, M. R., Campana, S., Covino, S., et al. 2003, A&A 399, 351
- Paturel, G., Andernach, H., Bottinelli, L., et al. 1997, A&AS 124, 109
- Paturel, G., Fouqué, P., Bottinelli, L. & Gougouenheim, L. 1989, A&AS 80 299
- Pildis, R. A., Bregman, J. N. & Evrard, A. E. 1995, ApJ 443, 514
- Ponman, T. J., Bourner, P. D. J., Ebeling, H., & Böhringer, H. 1996, MNRAS 283, 690
- Ranalli, P., Comastri, A. & Setti, G., proc. conf. *Multiwavelength AGN Surveys*, Cozumel, Mexico 2003, astro-ph/0404087
- Read, A. M., Ponman, T. J. & Strickland, D. K. 1997, MNRAS 286, 626
- Roberts, T. P. & Warwick, R. S. 2000, MNRAS 315, 98
- Sandage, A. & Tammann, G. A., *Revised Shapley-Ames Catalog of Bright Galaxies* 1981, Carnegie Inst. of Washington, Publ. 635
- Severgnini, P., Caccianiga, A., Braitto, V., et al. 2003, A&A, 406, 483
- Shapley, A., Fabbiano, G. & Eskridge, P. B. 2001, ApJS 137, 139
- Stocke, J. T., Morris, S. L., Gioia, I. M., et al. 1991, ApJS 76, 813
- Trinchieri, G., Noris, L. & di Serego Alighieri, S., 1997, A&A 326, 565
- Trinchieri, G., Fabbiano, G. & Peres, G. 1988, ApJ 325, 531
- Tully, R. B., *Nearby galaxies catalog*, Cambridge and New York, Cambridge University Press, 1988
- de Vaucouleurs, G., de Vaucouleurs, A. & Corwin, J. R., *Second reference catalogue of bright galaxies* 1976, Austin: University of Texas Press
- Véron-Cetty, M.-P. & Véron, P., 2001, A&A 374, 92
- Voges, W., Aschenbach, B., Boller, T., et al. 1999, A&A 349, 389
- Vogler, A. & Pietsch, W. 1996, A&A, 311, 35
- White, N. E., Giommi, P., & Angelini, L. 1994, Bulletin of the American Astronomical Society, 26, 1372
- Zimmermann, H.-U., Boller, T., Döbereiner, S. & Pietsch, W. 2001, A&A 378, 30




Fig. 2. First page of the figure available in electronic form, that presents the X-ray contours from smoothed X-ray images superimposed onto optical images. Galaxies are generally at the center of the field. Smoothing is done with a Gaussian function of $\sigma = 5''$ for point-like sources and of $\sigma = 10''$ for extended sources.

Appendix A: Results of optical observations

The log of observations made at the Loiano Telescope in order to measure redshifts and magnitudes for some galaxies in our sample (see section 5) is listed in Table A.1.

The seeing was about $2''$, so we used a long slit of width $2.5''$ for spectra. The spectral range was $3800 \text{ \AA} - 8700 \text{ \AA}$.

For photometry we used the B - V - R filters of Johnson - Kron - Cousin system.

We made use of the Image Reduction and Analysis Facility (IRAF)⁶ for spectroscopic and photometric data reduction.

We used all lines visible in the spectra to measure z , which is calculated from the average value. The redshifts obtained are reported in Table A.2. Only absorption lines, typical of early-type galaxies, and no sign of nuclear activity were detected in the galaxy spectra.

The atmospheric conditions were inadequate for photometric observations, however we list our raw estimate of magnitudes in Table A.3 and we use the value obtained for PGC 69291 in Table 1, where it is reported in brackets.

PGC 196693

After obtaining the spectrum, we discovered that NED lists the galaxy SDSS J234817.99+010617.0 at the same position and with the same redshift we obtained, although it does not give a PGC name. We therefore identify PGC 196693 as SDSS J234817.99+010617.0.

PGC 69291

The galaxy is identified as NGC 7325 by LEDA and as NGC 7327 by NED, with a note suggesting that it could be a star. Our observations show 2 objects, a star and a galaxy at $z = 0.0290 \pm 0.0009$ at the position of PGC 69291. Most of the optical flux is due to the star, as listed in Table A.3.

The uncertainty in the X-ray position (see also the map in Fig. 2) does not allow us to identify the X-ray source with either object. Unfortunately the bad quality of our optical does not allow us to use the X-ray - to - optical ratio (see for example Maccacaro et al., 1988) to discriminate between them, because it is consistent with both the star and the galaxy. We list the galaxy in the total sample but due to its faint estimated magnitude it is not included in the complete serendipitous sample.

Appendix B: Table of total sample

We report here the complete table with our total sample that will be available in the electronic form. For explanation of the columns refer to the sample Table 1 in the main text.

⁶ IRAF is a general purpose software system for the reduction and analysis of astronomical data. IRAF is written and supported by the IRAF programming group at the National Optical Astronomy Observatories (NOAO) in Tucson, Arizona. NOAO is operated by the Association of Universities for Research in Astronomy (AURA), Inc. under cooperative agreement with the National Science Foundation.

Name	LEDA coordinates		Spectroscopy Exposure (s)	Photometry	
	R. A. (2000)	Dec. (2000)		Filter	Exposure (s)
PGC 65435	20 48 43.09	25 16 53.0	2×1800	R	2×120
PGC 69291	22 36 33.58	34 30 04.5	2×1800	R	1×60
				R	3×40
				V	1×40
				V	3×80
				B	3×200
PGC 196693	23 48 18.07	01 06 17.3	1×1800		
Feige 110	23 19 58.40	-05 09 56.2	2×480	R	2×30
				V	2×30
				B	2×40

Table A.1. Log of optical observations.

Name	Redshift
PGC 65435	0.0485 ± 0.0006
PGC 69291	0.0290 ± 0.0009
PGC 196693	0.093 ± 0.001

Table A.2. Average value of redshifts measured for the three galaxies observed.

Name	Magnitude		
	B	V	R
PGC 65435			14.8
PGC 69291 galaxy	16.6	16.1	15.6
PGC 69291 star	13.0	12.5	12.2

Table A.3. Estimated magnitudes for two of the three galaxies observed and for the star found near PGC 69291 resulting from our observations with the Loiano Telescope.

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	
BMW000523.9+161307	00	05	23.83	+16	13	11.3	e	52	PGC 0000372		698		15.74	2.49	7.40	1.17	43.59	C1
BMW002055.1+215208	00	20	55.43	+21	51	52.2	p		PGC 0001333 IC 1543	Sbc	112	14.54	7.63	1.11	3.58	0.52	41.71	c
BMW002055.2+215208																		
BMW002549.3-453227	00	25	49.26	-45	32	26.5	p		PGC 0143535	Sab		16.57	18.64	3.16	7.64	1.30		
BMW002950.2-405630	00	29	50.99	-40	56	37.9	p		PGC 0130966 DUKST 294-9	Sc	241	15.35	19.06	2.31	7.81	0.95	42.72	c
BMW003652.3-333310	00	36	52.82	-33	33	14.7	p		PGC 0002204 ESO 350-IG38	S?	123	14.96	2.74	0.43	1.12	0.14	41.27	c
BMW003918.5+030220	00	39	18.47	+03	02	14.8	p		PGC 0002362 NGC 194	E	103	13.09	14.75	1.94	6.49	0.85	41.89	c; Gr
BMW003948.3+032219	00	39	48.77	+03	22	21.0	p		PGC 0002401 UM 57				6.15	0.49	2.70	0.22		
BMW004242.0+405154	00	42	42.48	+40	51	52.6	e	60	PGC 0002555 NGC 221	E	1	8.18	52.44	1.06	26.75	0.54	38.52	Gr
BMW010716.2+323117	01	07	16.20	+32	31	16.6	p		PGC 0003966 NGC 379	S0	110	13.48	1.37	0.35	0.67	0.17	40.98	Gr
BMW010717.4+322857	01	07	17.76	+32	28	57.0	p		PGC 0003969 NGC 380	E	88	13.24	4.19	0.50	2.06	0.24	41.26	Gr
BMW010717.8+322857																		
BMW010918.1+131013	01	09	18.63	+13	10	11.4	e	60	PGC 0004117 UGC 716		356	14.78	3.77	0.67	1.77	0.32	42.41	c; C1
BMW011314.7+153059	01	13	15.84	+15	31	05.0	p		PGC 0004392		276		1.60	0.24	0.75	0.12	41.78	C1
BMW011315.3+153102																		
BMW011315.8+153059																		
BMW012038.2+294157	01	20	38.06	+29	41	56.2	p		PGC 0004848 IC 1672	Sb	142	13.28	7.83	2.65	4.00	1.35	41.96	c
BMW012038.8+294150																		
BMW012310.6+332738	01	23	11.33	+33	27	29.5	e	50	PGC 0005060 NGC 499	E-S0	87	12.86	28.37	1.51	13.90	0.74	42.01	Gr
BMW012311.5+332741																		
BMW012558.8-012023	01	25	59.10	-01	20	22.8	e	18	PGC 0005323 NGC 545	E-S0	107	12.78	1.72	0.28	0.81	0.13	41.02	c; C1
BMW012600.2-012040	01	26	00.44	-01	20	40.3	e	50	PGC 0005324 NGC 547	E	110	12.68	4.83	0.69	2.27	0.32	41.50	c; C1
BMW012600.4-012039																		
BMW012600.7-012040																		
BMW012623.1+013921	01	26	23.24	+01	39	17.7	p		PGC 0144123 APMUKS(BJ)B012348.87+012346.1			17.13	9.92	0.84	4.37	0.37		
BMW014358.8+022053	01	43	58.82	+02	20	52.5	e	40	PGC 0006367 UGC 1214	S0-a	103	14.49	15.22	0.77	6.70	0.34	41.90	c
BMW015641.6+330238	01	56	41.61	+33	02	30.9	p		PGC 0007289 NGC 736	E	87	12.72	1.17	0.33	0.60	0.17	40.71	c; Gr
BMW021853.5+423440	02	18	53.93	+42	34	38.9	p		PGC 0165335	E	110		0.48	0.14	0.26	0.08	40.53	C1
BMW022537.8+365759	02	25	37.90	+36	57	48.7	p		PGC 0009215 UGC 1877	E	216	14.14	12.86	1.02	6.30	0.50	42.52	C1
BMW022602.5-232136	02	26	02.54	-23	21	36.0	p		PGC 0133905 DUKST 479-24	Sc	337	15.59	2.79	0.07	1.14	0.03	42.15	
BMW023741.8+015827	02	37	41.49	+01	58	27.5	p		PGC 0009961 NGC 1004	E	130	13.70	5.19	1.18	1.84	0.52	41.58	C1
BMW023819.3+020706	02	38	19.35	+02	07	12.5	p		PGC 0009997 NGC 1016	E	132	12.59	3.58	1.16	1.58	0.51	41.47	C1
BMW024539.7-525748	02	45	39.70	-52	57	47.5	e	40	PGC 0010446 \protect\vrule width0pt\protect\href{http://arXiv.org/abs/eso-lv/1540110}{eso-lv 1540110}									
BMW031858.7+412808	03	18	58.42	+41	28	08.3	p		PGC 0012350 NGC 1270	E	98	13.40	1.61	0.31	0.93	0.18	40.99	C1
BMW032244.0-370614	03	22	44.13	-37	06	12.2	p		PGC 0012653 NGC 1317	SBa	25	11.58	3.27	0.36	1.34	0.15	40.00	C1
BMW032244.1-370613																		
BMW033051.3-523034	03	30	51.46	-52	30	36.3	p		PGC 0013046	E-S0	345	14.98	2.21	0.89	0.87	0.36	42.08	C1
BMW033657.5-353015	03	36	57.51	-35	30	15.1	p		PGC 0013344 NGC 1387	E-S0	25	11.69	6.76	0.59	2.50	0.22	40.26	C1
BMW033728.0-243003	03	37	28.17	-24	30	05.8	p		PGC 0013368 NGC 1385	SBc	26	10.87	6.20	1.59	2.29	0.59	40.26	c
BMW033828.6-352659	03	38	28.90	-35	27	02.1	e	500	PGC 0013418 NGC 1399	E	25	10.06	411.10	22.67	152.10	8.39	42.03	C1
BMW033828.9-352659																		
BMW033830.0-352711																		
BMW033851.5-353536	03	38	51.92	-35	35	41.9	e	150	PGC 0013433 NGC 1404	E	25	10.62	111.00	1.21	41.06	0.45	41.46	C1
BMW033851.6-353536																		
BMW033851.7-353536																		
BMW041205.3-325222	04	12	03.09	-32	52	28.9	e	75 150	PGC 0014638 NGC 1532	SBb	20	9.69	5.16	1.26	2.12	0.52	40.00	Gr
BMW041544.5-553533	04	15	45.64	-55	35	33.2	e	60	PGC 0014757 NGC 1549	E	20	10.36	7.72	0.78	3.17	0.32	40.16	Gr
BMW041545.1-553537																		
BMW041545.8-553533																		

BMW042900.6-533653	04	29	00.91	-53	36	45.0	p	PGC 0015231	IC 2081	E-S0	255	14.62	3.39	0.60	1.25	0.22	41.98	C1
BMW042900.7-533647																		
BMW043754.0-425852	04	37	55.01	-42	58	53.0	p	PGC 0130376		E		15.53	12.38	2.04	5.08	0.84		
BMW044707.0-202847	04	47	06.29	-20	28	39.2	p	PGC 0099957		E	430	15.41	6.05	1.63	2.66	0.72	42.75	C1
BMW050138.5-041527	05	01	38.26	-04	15	33.5	p	PGC 0016574	NGC 1741	Sm	81	14.56	3.60	0.91	1.84	0.46	41.15	c; Gr
BMW050943.5-083652	05	09	44.01	-08	36	51.9	p	PGC 0147040		S0-a		17.00	0.89	0.34	0.49	0.19		
BMW053359.9-714524	05	34	00.41	-71	45	28.5	p	PGC 0017451	ESO 56-G154		145	13.42	16.17	1.94	8.90	1.07	42.33	c
BMW062457.1+821914	06	24	57.12	+82	19	13.8	p	PGC 0018991	UGC 3435		86	14.19	3.17	0.82	1.55	0.40	41.10	c
BMW062545.7-042202	06	25	45.47	-04	22	04.4	p	PGC 0075751	CGMW 1-0108	Sc			12.25	0.66	11.39	0.62		
BMW062545.8-042203																		
BMW071957.5-320041	07	19	58.78	-32	00	54.4	p	PGC 0077028	CGMW 2-0778	Sbc		13.74	17.30	3.91	10.03	2.27		
BMW073217.8+854234	07	32	17.80	+85	42	34.1	e	PGC 0021231	NGC 2300	E-S0	47	11.61	10.90	0.43	5.56	0.22	41.15	Gr
BMW073220.2+854230																		
BMW074914.6-055607	07	49	14.56	-05	56	11.3	p	PGC 0078533	CGMW 1-1742	Sc			1.25	0.26	0.72	0.15		
BMW081023.2+421628	08	10	23.16	+42	16	27.1	e	PGC 0022928	CGCG 207-40		379	15.90	22.15	1.17	10.85	0.57	43.25	C1
BMW082133.8+470242	08	21	33.55	+47	02	37.7	p	PGC 0023450			775	17.61	1.27	0.59	0.60	0.28	42.61	C1
BMW090825.4-093349	09	08	25.49	-09	33	41.9	p	PGC 0025701			958	17.27	1.69	0.26	0.82	0.13	42.92	C1
BMW090825.7-093336																		
BMW090851.5+110122	09	08	51.50	+11	01	22.0	e	PGC 0025741			978	17.07	1.50	0.31	0.66	0.14	42.88	C1
BMW090852.1+110138																		
BMW090943.9+071032	09	09	43.87	+07	10	31.5	p	PGC 0025825	NGC 2773	Sb	109	14.26	1.05	0.32	0.50	0.15	40.83	c
BMW091041.7+071223	09	10	41.86	+07	12	27.0	p	PGC 0025876	NGC 2777	Sab	26	14.45	1.67	0.31	0.79	0.15	39.79	Gr
BMW091826.3-122222	09	18	27.00	-12	22	26.9	e	PGC 0087445			315	15.68	9.97	1.13	4.89	0.56	42.74	C1
BMW100108.4-192631	10	01	08.92	-19	26	26.8	p	PGC 0028995	ESO 567-G3	Sab		14.30	10.91	1.26	5.34	0.61		c; Gr?
BMW100108.7-192621																		
BMW100114.9+554316	10	01	15.17	+55	43	06.0	p	PGC 0028990	MCG 9-17-9		31	15.20	1.10	0.38	0.41	0.14	39.64	c; Gr
BMW100115.1+554318																		
BMW100115.5+554308																		
BMW100158.3+554100	10	01	58.60	+55	40	49.0	e	PGC 0029050	NGC 3079	SBcd	31	9.84	18.44	1.18	6.82	0.44	40.87	c; Gr
BMW100158.3+554060																		
BMW100158.4+554103																		
BMW100158.7+554051																		
BMW101010.1+541828	10	10	10.10	+54	18	27.7	p	PGC 0029594	CGCG 266-27		261	15.38	2.57	1.27	0.95	0.47	41.84	
BMW101825.0+215348	10	18	25.03	+21	53	45.9	e	PGC 0030099	NGC 3193	E	35	11.72	1.21	0.31	0.50	0.13	39.84	Gr
BMW103633.4-271318	10	36	33.36	-27	13	17.4	e	PGC 0031453			215	14.71	9.12	2.81	4.47	1.38	42.42	C1
BMW103719.0-271121	10	37	18.92	-27	11	24.8	e	PGC 0031540	NGC 3315	E-S0	75	13.84	1.39	0.36	0.68	0.18	40.65	C1
BMW104827.8+123157	10	48	27.76	+12	31	56.6	e	PGC 0032306	NGC 3389	Sc	34	11.63	3.23	0.89	1.42	0.39	40.26	Gr
BMW104925.3+324639	10	49	25.13	+32	46	27.1	p	PGC 0032390	IC 2604	SBm	50	14.18	2.45	0.51	1.01	0.21	40.45	
BMW110522.3+381421	11	05	23.18	+38	14	13.2	p	PGC 0033514	CGCG 213-27		170	15.92	9.06	2.70	3.35	1.00	42.05	
BMW113446.5+485718	11	34	46.51	+48	57	17.6	p	PGC 0035780	IC 711	E-S0	194	14.95	1.39	0.41	0.57	0.17	41.41	C1
BMW113448.9+490440	11	34	49.51	+49	04	39.6	p	PGC 0035785	IC 712	E	201	13.67	1.48	0.45	0.61	0.18	41.44	C1
BMW114223.6+101552	11	42	23.79	+10	15	51.2	p	PGC 0036348	NGC 3825	SBa	128	13.57	1.26	0.33	0.55	0.15	40.97	Gr
BMW114232.1+583706	11	42	32.04	+58	37	05.8	p	PGC 0036369	MCG +10-17-46			15.82	1.21	0.32	0.45	0.12		
BMW114430.3+305305	11	44	30.20	+30	53	02.9	p	PGC 0139685			458	16.51	4.29	1.43	1.76	0.59	42.65	C1
BMW115535.8+232724	11	55	35.92	+23	27	27.5	p	PGC 0139725			815	17.55	3.24	0.57	1.33	0.23	42.99	
BMW120438.2+014717	12	04	38.50	+01	47	10.5	p	PGC 0038218	NGC 4077	S0	140	13.97	1.68	0.64	0.69	0.26	41.19	Gr
BMW120631.5-295119	12	06	31.46	-29	51	18.7	p	PGC 0157406				14.62	0.52	0.17	0.25	0.08		c
BMW121049.5+392822	12	10	49.82	+39	28	23.9	p	PGC 0038773	NGC 4156	SBb	135	13.65	1.71	0.13	0.70	0.05	41.17	c
BMW121049.5+392820																		
BMW121049.6+392823																		
BMW121633.3+072746	12	16	34.07	+07	27	46.9	p	PGC 0039328	NGC 4224	Sa	53	12.35	5.79	1.06	2.37	0.43	40.88	C1

BMW121908.2+470531	12	19	08.40	+47	05	20.5	p		PGC 0039615	UGC 7356	Irr	8	15.35	1.87	0.47	0.96	0.17	38.65			
BMW122018.0+752214	12	20	17.39	+75	22	16.6	e	40	PGC 0039791	NGC 4291	E	44	12.04	13.65	1.37	6.01	0.60	41.11			
BMW122503.5+125317	12	25	04.23	+12	53	07.5	e	100	PGC 0040455	NGC 4374	E	25	9.84	54.48	2.77	23.97	1.22	41.23	Cl		
BMW122504.1+125308																					
BMW122504.3+125318																					
BMW122504.4+125316																					
BMW122504.4+125319																					
BMW122611.1+125642	12	26	11.68	+12	56	45.2	e		200	400	PGC 0040653	NGC 4406	E	25	9.73	199.80	7.89	87.91	3.47	41.79	Cl
BMW122612.6+125628																					
BMW122716.2+085815	12	27	14.97	+08	58	17.3	e	80	PGC 0094216			530	16.11	3.74	1.03	1.53	0.42	42.73	Cl		
BMW123017.6+121936	12	30	17.59	+12	19	35.7	e	30	PGC 0041297	NGC 4478	E	25	11.93	1.31	0.30	0.68	0.13	39.63	Cl		
BMW123018.1+121926																					
BMW123018.1+121926																					
BMW123052.5+143307	12	30	52.51	+14	33	06.7	e	40	PGC 0169455	VPC 774			18.18	7.98	1.43	3.51	0.63				
BMW123643.2+131515	12	36	43.14	+13	15	19.0	p		PGC 0042081	IC 3583	Irr	25	13.11	0.67	0.32	0.30	0.14	39.33	Cl		
BMW124342.7+162340	12	43	42.68	+16	23	38.0	p		PGC 0042833	NGC 4651	Sc	25	10.81	1.30	0.36	0.53	0.15	39.55	c; Cl		
BMW125232.8-153057	12	52	33.03	-15	31	02.2	p		PGC 0043675	MCG -2-33-38	S0	87	14.39	3.19	0.31	1.50	0.14	41.11	c; Gr		
BMW125233.1-153102																					
BMW125302.3-152120	12	53	02.26	-15	21	19.7	p		PGC 0043748	DRCG25-68	E-S0	266	15.55	0.54	0.20	0.25	0.09	41.38	Cl		
BMW125318.1-153203	12	53	18.27	-15	32	03.4	e	30	PGC 0043777	DRCG25-39	E-S0	284	14.78	2.94	0.34	1.38	0.16	42.11	Cl		
BMW125318.3-153203																					
BMW125749.0-171628	12	57	49.85	-17	16	25.3	e	50	PGC 0044369		E	296	14.77	13.80	1.77	6.76	0.87	42.84	Cl		
BMW125750.2-171616																					
BMW130055.7+274740	13	00	55.99	+27	47	29.8	p		PGC 0044840	NGC 4911	SBbc	159	13.30	12.83	1.03	4.74	0.38	42.14	Cl		
BMW130056.0+274737																					
BMW130056.2+274731																					
BMW130559.7+291647	13	05	59.46	+29	16	46.8	p		PGC 0045318	CGCG 160-136	E	159	15.08	4.42	0.95	1.64	0.35	41.66	c		
BMW130803.0+292918	13	08	03.09	+29	29	12.0	p		PGC 0166974					2.61	0.17	0.97	0.06		Cl?		
BMW130803.2+292910																					
BMW130803.3+292911																					
BMW130803.4+292903																					
BMW131324.5-325247	13	13	24.96	-32	52	44.7	p		PGC 0158561				16.26	1.12	0.41	0.55	0.20				
BMW131413.5-425739	13	14	13.44	-42	57	38.8	e	60	PGC 0046023	NGC 5026	SBa	75	11.73	86.49	3.71	47.57	2.04	42.48	Cl?		
BMW131912.4-123228	13	19	12.40	-12	32	28.1	e	20	PGC 0046432	NGC 5072	E-S0	136	14.23	2.09	0.74	0.92	0.33	41.27	c		
BMW132040.2-434200	13	20	40.20	-43	42	00.3	p		PGC 0046566	NGC 5082	S0	78	12.96	0.93	0.50	0.51	0.27	40.54	Gr		
BMW132745.6-314751	13	27	45.41	-31	47	54.0	p		PGC 0047177			265	14.53	3.74	0.61	1.76	0.29	42.15	Cl		
BMW132759.6-272104	13	27	59.87	-27	21	29.6	p		PGC 0047209	IC 4255	E	203	13.66	3.91	1.11	1.92	0.55	41.95	Cl		
BMW132802.6-314519	13	28	02.56	-31	45	19.0	p		PGC 0088857		E	258	15.01	3.37	0.60	1.58	0.28	42.08	Cl		
BMW132958.2+471613	13	30	00.20	+47	15	55.1	e	80	PGC 0047413	NGC 5195	SBpec	14	10.10	12.66	0.77	5.19	0.32	40.06	Gr		
BMW132959.1+471555																					
BMW132959.8+471556																					
BMW133000.1+471558																					
BMW133006.4-014316	13	30	06.34	-01	43	06.8	e	45	PGC 0047432	NGC 5183	Sb	86	13.21	2.86	0.98	1.17	0.40	41.04	c		
BMW133425.1+344127	13	34	24.86	+34	41	26.9	e	60	PGC 0047822	NGC 5223	E	144	13.22	20.33	1.70	7.52	0.63	42.24	Cl		
BMW134623.6-375810	13	46	23.36	-37	58	23.3	p		PGC 0048817	ESO 325-G16		228	14.67	11.43	1.27	5.37	0.60	42.48	Cl		
BMW135321.7+402151	13	53	21.67	+40	21	50.7	p		PGC 0049347	NGC 5350	SBbc	57	11.87	1.38	0.40	0.51	0.15	40.26	Gr		
BMW140715.2-270922	14	07	15.27	-27	09	27.7	e	30	PGC 0050373	ESO 510-G66	E-S0	146	13.97	9.04	0.77	4.43	0.38	42.03	Cl		
BMW141319.7-030857	14	13	19.76	-03	08	58.0	p		PGC 0050786	NGC 5507	S0	43	13.22	0.31	0.09	0.14	0.04	39.47			
BMW144104.0+532011	14	41	04.03	+53	20	10.7	p		PGC 0084264			629	17.53	5.88	1.02	2.18	0.38	42.99			
BMW151643.2+552432	15	16	42.26	+55	24	22.2	p		PGC 0054522	NGC 5908	Sb	66	12.20	0.53	0.20	0.20	0.07	39.92			
BMW160434.7+174315	16	04	34.65	+17	43	21.9	e	300	PGC 0056962	NGC 6041	E-S0	209	13.90	183.30	4.47	80.65	1.97	43.60	Cl		

[illegible]